

Continuous Distributions

A normal distribution and other density functions involving exponential forms play the most important role in probability and statistics. They are related in a certain way, as summarized in a diagram later in this topic.

Exponential distribution. The exponential density function is defined as

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & x \geq 0; \\ 0 & x < 0 \end{cases}$$

Then the cdf is computed as

$$F(t) = \begin{cases} 1 - e^{-\lambda t} & t \geq 0; \\ 0 & t < 0 \end{cases}$$

where $\lambda > 0$ is called a **rate parameter**.

Example 1. Suppose that the number of wins in a slot machine for an hour is distributed with Poisson distribution with λ . (a) If you play for t hours, what is the distribution for the number of wins in t hours? (b) Let X be the exact time of your first win. Find $P(X > t)$. (c) What is the distribution for X ?

Solution. (a) It is again a Poisson distribution with parameter λt . (b) Let Y be the number of wins in t hours. Then we have $P(X > t) = P(Y = 0)$. Thus,

$$P(X > t) = e^{-\lambda t} \frac{\lambda^0}{0!} = e^{-\lambda t}$$

(c) Since $F(t) = P(X \leq t) = 1 - P(X > t) = 1 - e^{-\lambda t}$, X must be exponentially distributed.

Survival function and memoryless property. The function $S(t) = 1 - F(t) = P(X > t)$ is called the **survival function**. When $F(t)$ is an exponential distribution, we have $S(t) = e^{-\lambda t}$ for $t \geq 0$. Furthermore, we can find that

$$P(X > t + s | X > s) = \frac{P(X > t + s)}{P(X > s)} = \frac{S(t + s)}{S(s)} = S(t) = P(X > t), \quad (4.1)$$

which is referred as the **memoryless** property of exponential distribution.

Example 2. A service time (in minutes) of post office is exponentially distributed with $\lambda = 0.2$. (a) Suppose that no one is there, and that you get the service immediately. Find the probability that it takes more than 5 minutes to complete the business. (b) Suppose that someone else is in service ahead of you, and that you are next in line. Find the probability that you wait more than 5 minutes.

Solution. Let X be the service time. (a) We obtain

$$P(X > 5) = 1 - P(X \leq 5) = 1 - F(5) = e^{-1} \approx 0.37.$$

(b) The service of the person ahead of you started s minutes ago. $P(X > 5 + s | X > s) = P(X > 5) \approx 0.37$.

Gamma distribution. The *gamma function*, denoted by $\Gamma(\alpha)$, is defined as

$$\Gamma(\alpha) = \int_0^{\infty} u^{\alpha-1} e^{-u} du, \quad \alpha > 0.$$

Then the *gamma density* is defined as

$$f(t) = \frac{\lambda^\alpha}{\Gamma(\alpha)} t^{\alpha-1} e^{-\lambda t} \quad t \geq 0$$

which depends on two parameters $\alpha > 0$ and $\lambda > 0$. [The textbook uses a parameter $\beta = \frac{1}{\lambda}$ instead.]

Properties of gamma function.

$$\begin{aligned} \Gamma(1) &= \int_0^{\infty} e^{-u} du = 1 \\ \Gamma(\alpha) &= \int_0^{\infty} u^{\alpha-1} e^{-u} du \\ &= [-e^{-u} u^{\alpha-1}]_{u=0}^{u=\infty} + (\alpha-1) \int_0^{\infty} u^{\alpha-2} e^{-u} du \\ &= (\alpha-1) \Gamma(\alpha-1) \end{aligned}$$

Suppose that the parameter α is a positive integer n . Then we have

$$\begin{aligned} \Gamma(n) &= (n-1) \Gamma(n-1) = \dots \\ &= (n-1) \times \dots \times 2 \times 1 \times \Gamma(1) = (n-1)! \end{aligned}$$

The mgf of gamma distribution. Let X be a gamma random variable with parameter (α, λ) . For $t < \lambda$, the mgf $M_X(t)$ of X can be computed as follows.

$$\begin{aligned} M_X(t) &= \int_0^{\infty} e^{tx} \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} dx = \frac{\lambda^\alpha}{\Gamma(\alpha)} \int_0^{\infty} x^{\alpha-1} e^{-(\lambda-t)x} dx \\ &= \frac{\lambda^\alpha}{(\lambda-t)^\alpha} \frac{1}{\Gamma(\alpha)} \int_0^{\infty} ((\lambda-t)x)^{\alpha-1} e^{-(\lambda-t)x} (\lambda-t) dx \\ &= \left(\frac{\lambda}{\lambda-t} \right)^\alpha \frac{1}{\Gamma(\alpha)} \int_0^{\infty} u^{\alpha-1} e^{-u} du = \left(\frac{\lambda}{\lambda-t} \right)^\alpha \end{aligned}$$

If you use $\lambda = 1/\beta$ then we obtain $M_X(t) = (1 - \beta t)^{-\alpha}$.

Parameters of gamma/exponential distribution. We call the parameter α a **shape parameter**, because changing α changes the shape of the density. We call the parameter λ a **rate parameter**, because changing λ merely rescales the density without changing its shape. In particular, the gamma distribution with $\alpha = 1$ becomes an exponential distribution (with parameter λ).

The parameter $\beta = 1/\lambda$, if you choose, is called a **scale parameter**.

Property of scale parameter. Let X be a service time (in minutes) having a gamma distribution. If we change the unit of measurement from minutes to seconds then $(60)X$ has a gamma distribution with new scale parameter $(60)\beta$. While the scale parameter changes from β to $(60)\beta$, the rate parameter changes from λ to $\lambda/60$.

Theorem 3. *If X has a gamma distribution with parameters α and $\beta = 1/\lambda$ then $Y = kX$ has a gamma distribution with parameters α and $k\beta$.*

Solution. We obtain

$$M_Y(t) = M_X(kt) = [1 - \beta(kt)]^{-\alpha} = [1 - (k\beta)t]^{-\alpha}$$

Thus, Y has the mgf of gamma distribution with parameter α and $k\beta$.

Moments of gamma distribution. We can find the first and second moment

$$E[X] = M'_X(0) = \frac{\alpha}{\lambda}$$

and

$$E[X^2] = M''_X(0) = \frac{\alpha(\alpha + 1)}{\lambda^2}$$

Thus, we can obtain

$$\text{Var}(X) = \frac{\alpha(\alpha + 1)}{\lambda^2} - \frac{\alpha^2}{\lambda^2} = \frac{\alpha}{\lambda^2}.$$

If you use $\beta = 1/\lambda$ then $E[X] = \alpha\beta$ and $\text{Var}(X) = \alpha\beta^2$.

Expectation by direct calculation. Let X be a gamma random variable with parameter (α, λ) . Then we can calculate $E[X]$ and $E[X^2]$ as follows.

$$\begin{aligned} E[X] &= \int_0^{\infty} x \frac{\lambda^{\alpha}}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} dx = \frac{1}{\lambda \Gamma(\alpha)} \int_0^{\infty} (\lambda x)^{\alpha} e^{-\lambda x} \lambda dx \\ &= \frac{1}{\lambda \Gamma(\alpha)} \int_0^{\infty} u^{\alpha} e^{-u} du = \frac{1}{\lambda} \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha)} = \frac{\alpha}{\lambda} \end{aligned}$$

Variance by direct calculation. We have

$$\begin{aligned} E[X^2] &= \int_0^{\infty} x^2 \frac{\lambda^{\alpha}}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} dx = \frac{1}{\lambda^2 \Gamma(\alpha)} \int_0^{\infty} (\lambda x)^{\alpha+1} e^{-\lambda x} \lambda dx \\ &= \frac{1}{\lambda^2 \Gamma(\alpha)} \int_0^{\infty} u^{\alpha+1} e^{-u} du = \frac{1}{\lambda^2} \frac{\Gamma(\alpha + 2)}{\Gamma(\alpha)} = \frac{\alpha(\alpha + 1)}{\lambda^2} \end{aligned}$$

Thus, we can obtain

$$\text{Var}(X) = \frac{\alpha(\alpha + 1)}{\lambda^2} - \frac{\alpha^2}{\lambda^2} = \frac{\alpha}{\lambda^2}$$

Sum of independent random variables. Let X and Y be independent gamma random variables with the respective parameters (α_1, λ) and (α_2, λ) . Then the sum $Z = X + Y$ of random variables has the mgf

$$M_Z(t) = M_X(t) \cdot M_Y(t) = \left(\frac{\lambda}{\lambda - t}\right)^{\alpha_1} \cdot \left(\frac{\lambda}{\lambda - t}\right)^{\alpha_2} = \left(\frac{\lambda}{\lambda - t}\right)^{\alpha_1 + \alpha_2}$$

which is the mgf of gamma distribution with parameter $(\alpha_1 + \alpha_2, \lambda)$. Thus, Z is a gamma random variable with parameter $(\alpha_1 + \alpha_2, \lambda)$.

Chi-square distribution. The gamma distribution

$$f(x) = \frac{1}{2^{n/2}\Gamma(n/2)} x^{n/2-1} e^{-x/2}, \quad x \geq 0$$

with $\alpha = \frac{n}{2}$ and $\lambda = \frac{1}{2}$ is called the **chi-square distribution** with n degrees of freedom. [It plays a vital role later in understanding another important distribution, called t -distribution later.] A chi-square random variable X has the mean $E[X] = n$ and the variance $\text{Var}(X) = 2n$.

Example 4. Let X be a chi-square random variable with $n = 6$.

- (a) Find the quantile $Q(0.05)$ and $Q(0.95)$.
- (b) Let Y be a gamma random variable with $\alpha = 3$ and $\lambda = 1/4$. Then find $P(3.28 < Y < 25.2)$.

Solution.

- (a) From the table we obtain $Q(0.05) \approx 1.635$ and $Q(0.95) \approx 12.592$.
- (b) Since $X = Y/2$ has a chi-square random variable with $n = 6$, we obtain

$$\begin{aligned} P(3.28 < Y < 25.2) &= P(1.64 < X < 12.6) = F(12.6) - F(1.64) \\ &\approx 0.95 - 0.05 = 0.9 \end{aligned}$$

Normal distribution. A **normal distribution** is represented by a family of distributions which have the same general shape, sometimes described as “bell shaped.” The normal distribution has the pdf

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left[-\frac{(x - \mu)^2}{2\sigma^2}\right], \quad (4.2)$$

which depends upon two parameters μ and σ^2 . In (4.2), $\pi = 3.14159\dots$ is the famous “pi” (the ratio of the circumference of a circle to its diameter), and $\exp(u)$ is the exponential function e^u with the base $e = 2.71828\dots$ of the natural logarithm.

Integrations of normal density functions. Much celebrated integrations are the following:

$$\begin{aligned} \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx &= 1, \\ \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} x e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx &= \mu, \\ \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} (x-\mu)^2 e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx &= \sigma^2. \end{aligned} \tag{4.3}$$

Equation (4.3) guarantees that the function (4.2) always represents a “probability density” no matter what values the parameters μ and σ^2 would take.

Parameters of normal distribution. We say that a random variable X is **normally distributed** with parameter (μ, σ^2) when X has the pdf (4.2). Since $E[X] = \mu$ and $\text{Var}(X) = \sigma^2$, the parameter μ is called a **mean** (or, location parameter), and the parameter σ a **standard deviation** (or, a scale parameter). The parameter μ provides the center of distribution, and the density function $f(x)$ is symmetric around μ . Small values of σ lead to high peaks but sharp drops, while larger values of σ lead to flatter densities. The shorthand notation

$$X \sim N(\mu, \sigma^2)$$

is often used to express that X is a normal random variable with parameter (μ, σ^2) with **variance** σ^2 .

Linear transformation of normal distribution. One of the important properties of normal distribution is that their linear transformation remains normal.

Theorem 5. *Assume $a \neq 0$. If X is a normal random variable with parameter (μ, σ^2) [that is, the pdf of X is given by (4.2)], then $Y = aX + b$ is also a normal random variable having parameter $(a\mu + b, (a\sigma)^2)$.*

In particular,

$$\frac{X - \mu}{\sigma} \tag{4.4}$$

becomes a normal random variable with parameter $(0, 1)$, called the **standard normal distribution**.

Proof of linear transformation theorem. Assume that $a > 0$. Since X has the pdf (4.2), we obtain the cdf of Y as follows:

$$\begin{aligned} F_Y(t) &= P(aX + b \leq t) = P\left(X \leq \frac{t-b}{a}\right) \\ &= \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\frac{t-b}{a}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx \\ &= \frac{1}{a\sigma\sqrt{2\pi}} \int_{-\infty}^t e^{-\frac{(y-a\mu-b)^2}{2a^2\sigma^2}} dy \end{aligned}$$

Thus, Y has the pdf of $N(a\mu + b, a^2\sigma^2)$. The case with $a < 0$ is an exercise; see Problem 9.

Standard normal distribution. The normal density $\phi(x)$ with parameter $(0, 1)$ is given by

$$\phi(x) := \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}},$$

and the **table of standard normal distribution** is used to obtain the values for the cdf

$$\Phi(t) := \int_{-\infty}^t \phi(x) dx.$$

How to calculate probabilities. Suppose that a tomato plant height X is normally distributed with parameter (μ, σ^2) . Then what is the probability that the tomato plant height is between a and b ? The integration

$$P(a \leq X \leq b) = \int_a^b \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx$$

seems too complicated.

How to calculate probabilities, continued. If we consider the random variable $\frac{X - \mu}{\sigma}$, then the event $\{a \leq X \leq b\}$ is equivalent to the event

$$\left\{ \frac{a - \mu}{\sigma} \leq \frac{X - \mu}{\sigma} \leq \frac{b - \mu}{\sigma} \right\}.$$

Let $a' = \frac{a - \mu}{\sigma}$ and $b' = \frac{b - \mu}{\sigma}$. Then in terms of probability, this means that

$$P(a \leq X \leq b) = P\left(a' \leq \frac{X - \mu}{\sigma} \leq b'\right) = \int_{a'}^{b'} \phi(x) dx = \Phi(b') - \Phi(a'). \quad (4.5)$$

Finally look at the values for $\Phi(a')$ and $\Phi(b')$ from the table of standard normal distribution, and plug them into (4.5).

Example 6. The tomato plant height is normally distributed with parameter $(15, 16)$ in inches. What is the probability that the height is between 14.24 and 16.16 inches?

Solution. We can calculate $a' = \frac{14.24 - 15}{4} = -0.19$ and $b' = \frac{16.16 - 15}{4} = 0.29$. Then find $\Phi(-0.19) = 0.4247$ and $\Phi(0.29) = 0.6141$. Thus, the probability of interest becomes 0.1894, or approximately 0.19.

Summary of probability calculations. Suppose that a random variable X is normally distributed with mean μ and standard deviation σ . Then we can obtain

$$(a) P(a \leq X \leq b) = \Phi\left(\frac{b - \mu}{\sigma}\right) - \Phi\left(\frac{a - \mu}{\sigma}\right);$$

$$(b) P(X \leq b) = \Phi\left(\frac{b - \mu}{\sigma}\right);$$

$$(c) P(a \leq X) = 1 - \Phi\left(\frac{a - \mu}{\sigma}\right).$$

The mgf of normal distribution. Let X be a standard normal random variable. Then we can compute the mgf of X as follows.

$$\begin{aligned} M_X(t) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{tx} e^{-\frac{x^2}{2}} dx = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{1}{2}(x-t)^2 + \frac{t^2}{2}} dx \\ &= e^{\frac{t^2}{2}} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{(x-t)^2}{2}} dx = \exp\left(\frac{t^2}{2}\right) \end{aligned}$$

Since $X = (Y - \mu)/\sigma$ becomes a standard normal random variable and $Y = \sigma X + \mu$, the mgf $M(t)$ of Y can be given by

$$M(t) = e^{\mu t} M_X(\sigma t) = \exp\left(\frac{\sigma^2 t^2}{2} + \mu t\right).$$

Moments of normal distribution. Then we can find the first and second moment

$$E[X] = M'(0) = \mu \quad \text{and} \quad E[X^2] = M''(0) = \mu^2 + \sigma^2$$

Thus, we can obtain $\text{Var}(X) = (\mu^2 + \sigma^2) - \mu^2 = \sigma^2$.

Sum of independent random variables Let X and Y be independent normal random variables with the respective parameters (μ_x, σ_x^2) and (μ_y, σ_y^2) . Then the sum $Z = X + Y$ of random variables has the mgf

$$\begin{aligned} M_Z(t) &= M_X(t) \cdot M_Y(t) = \exp\left(\frac{\sigma_x^2 t^2}{2} + \mu_x t\right) \cdot \exp\left(\frac{\sigma_y^2 t^2}{2} + \mu_y t\right) \\ &= \exp\left(\frac{(\sigma_x^2 + \sigma_y^2)t^2}{2} + (\mu_x + \mu_y)t\right) \end{aligned}$$

which is the mgf of normal distribution with parameter $(\mu_x + \mu_y, \sigma_x^2 + \sigma_y^2)$. By the property (a) of mgf, we can find that Z is a normal random variable with parameter $(\mu_x + \mu_y, \sigma_x^2 + \sigma_y^2)$.

Relation between normal and chi-square distribution. Let X be a standard normal random variable, and let

$$Y = X^2$$

be the square of X . Then we can express the cdf F_Y of Y as $F_Y(t) = P(X^2 \leq t) = P(-\sqrt{t} \leq X \leq \sqrt{t}) = \Phi(\sqrt{t}) - \Phi(-\sqrt{t})$ in terms of the standard normal cdf Φ . By differentiating the cdf, we can obtain the pdf f_Y as follows.

$$f_Y(t) = \frac{d}{dt} F_Y(t) = \frac{1}{\sqrt{2\pi}} t^{-\frac{1}{2}} e^{-\frac{t}{2}} = \frac{1}{2^{1/2} \Gamma(1/2)} t^{-1/2} e^{-t/2},$$

where we use $\Gamma(1/2) = \sqrt{\pi}$. Thus, the square X^2 of X is the chi-square random variable with 1 degree of freedom.

The relation, continued. Now let X_1, \dots, X_n be independent and identically distributed random variables (iid random variables) of standard normal distribution. Since X_i^2 's have the gamma distribution with parameter $(1/2, 1/2)$, the sum

$$Y = \sum_{i=1}^n X_i^2$$


has the gamma distribution with parameter $(n/2, 1/2)$. That is, the sum Y has the chi-square distribution with n degree of freedom.

Extrema of exponential random variables. Let X_1, \dots, X_n be independent exponential random variables with the respective parameters $\lambda_1, \dots, \lambda_n$. Then we can define the random variable $V = \min(X_1, \dots, X_n)$ of the minimum of X_1, \dots, X_n . To find the distribution of V , consider the survival function $P(V > t)$ of V and calculate as follows.

$$\begin{aligned} P(V > t) &= P(X_1 > t, \dots, X_n > t) \\ &= P(X_1 > t) \times \dots \times P(X_n > t) \\ &= e^{-\lambda_1 t} \times \dots \times e^{-\lambda_n t} = e^{-(\lambda_1 + \dots + \lambda_n)t}, \end{aligned}$$

which is the survival function of exponential distribution with parameter $\lambda = \lambda_1 + \dots + \lambda_n$. Thus, V has an exponential distribution with parameter $\lambda = \lambda_1 + \dots + \lambda_n$.

Summary diagram.



Relationship among random variables. (1) If X_1 and X_2 are independent normal random variables with respective parameters (μ_1, σ_1^2) and (μ_2, σ_2^2) , then $X = X_1 + X_2$ is a normal random variable with $(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$.

(2) If X_1, \dots, X_m are iid normal random variables with parameter (μ, σ^2) , then $X = \sum_{k=1}^m \left(\frac{X_k - \mu}{\sigma} \right)^2$ is a chi-square (χ^2) random variable with m degrees of freedom.

(3) $\alpha = \frac{m}{2}$ and $\lambda = \frac{1}{2}$.

(4) If X_1, \dots, X_n are independent exponential random variables with respective parameters $\lambda_1, \dots, \lambda_n$, then $X = \min\{X_1, \dots, X_n\}$ is an exponential random variable with parameter $\sum_{i=1}^n \lambda_i$.

Relationship among random variables, continued. (5) If X_1, \dots, X_n are iid exponential random variables with parameter λ , then $X = \sum_{k=1}^n X_k$ is a gamma random variable with $(\alpha = n, \lambda)$.

(6) $\alpha = 1$.

(7) If X_1 and X_2 are independent random variables having gamma distributions with respective parameters (α_1, λ) and (α_2, λ) , then $X = X_1 + X_2$ is a gamma random variable with parameter $(\alpha_1 + \alpha_2, \lambda)$.

Exercises

Problem 1. Suppose that the lifetime of an electronic component follows an exponential distribution with rate parameter $\lambda = 0.2$.

- (a) Find the probability that the lifetime is less than 10.
- (b) Find the probability that the lifetime is between 5 and 15.

Problem 2. Suppose that in a certain population, individual's heights are approximately normally distributed with parameters $\mu = 70$ and $\sigma = 3$ in.

- (a) What proportion of the population is over 6ft. tall?
- (b) What is the distribution of heights if they are expressed in centimeters?

Problem 3. Let X be a normal random variable with $\mu = 5$ and $\sigma = 10$.

- (a) Find $P(X > 10)$.
- (b) Find $P(-20 < X < 15)$.
- (c) Find the value of x such that $P(X > x) = 0.05$.

Problem 4. Suppose that $X \sim N(\mu, \sigma^2)$.

- (a) Find $P(|X - \mu| \leq 0.675\sigma)$.
- (b) Find the value of c in terms of σ such that $P(\mu - c \leq X \leq \mu + c) = 0.95$.

Problem 5. X is an exponential random variable, and $P(X < 1) = 0.05$. What is λ ?

Problem 6. The lifetime in months of each component in a microchip is exponentially distributed with parameter $\mu = \frac{1}{10000}$.

- (a) For a particular component, we want to find the probability that the component breaks down before 5 months, and the probability that the component breaks down before 50 months. Use the approximation $e^{-x} \approx 1 - x$ for a small number x to calculate these probabilities.

- (b) Suppose that the microchip contains 1000 of such components and is operational until three of these components break down. The lifetimes of these components are independent and exponentially distributed with parameter $\mu = \frac{1}{10000}$. By using the Poisson approximation, find the probability that the microchip operates functionally for 5 months, and the probability that the microchip operates functionally for 50 months.

Problem 7. The length of a phone call can be represented by an exponential distribution. Then answer to the following questions.

- (a) A report says that 50% of phone call ends in 20 minutes. Assuming that this report is true, calculate the parameter λ for the exponential distribution.
- (b) Suppose that someone arrived immediately ahead of you at a public telephone booth. Using the result of (a), find the probability that you have to wait between 10 and 20 minutes.

Problem 8. Let $X_i, i = 1, \dots, 6$, be independent standard normal random variables. Find

$$P(X_1^2 + X_2^2 + X_3^2 + X_4^2 + X_5^2 + X_6^2 \leq 1.64)$$

Problem 9. Let X be a normal random variable with parameter (μ, σ^2) , and let $Y = aX + b$ with $a < 0$.

- (a) Show that

$$P(Y \leq t) = \int_{-\infty}^t f(x) dx,$$

where

$$f(x) = \frac{1}{|a|\sigma\sqrt{2\pi}} \exp \left[-\frac{(x - a\mu - b)^2}{2(a\sigma)^2} \right].$$

- (b) Conclude that Y is a normal random variable with parameter $(a\mu + b, (a\sigma)^2)$.

Problem 10. Let X be a normal random variable with parameter (μ, σ^2) , and let $Y = e^X$. Find the density function f of Y by forming

$$P(Y \leq t) = \int_0^t f(x) dx.$$

This density function f is called the *lognormal density* since $\ln Y$ is normally distributed.

Optional problems

Problem 11. Show that the standard normal density integrates to unity by completing the following questions.

(a) Show that
$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-\frac{x^2+y^2}{2}} dx dy = \int_0^{\infty} \int_0^{2\pi} e^{-\frac{r^2}{2}} r d\theta dr = 2\pi.$$

(b) Show that
$$\frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{y^2}{2}} dy = 1.$$

Problem 12. Suppose that a non-negative random variable X has the memoryless property

$$P(X > t + s \mid X > s) = P(X > t).$$

By completing the following questions, we will show that X *must* be an exponential random variable.

(a) Let $S(t)$ is the survival function of X . Show that the memoryless property implies that $S(s + t) = S(s)S(t)$ for $s, t \geq 0$.

(b) Let $\kappa = S(1)$. Show that $\kappa > 0$.

(c) Show that $S\left(\frac{1}{n}\right) = \kappa^{\frac{1}{n}}$ and $S\left(\frac{m}{n}\right) = \kappa^{\frac{m}{n}}$.

Therefore, we can obtain $S(t) = \kappa^t$ for any $t \geq 0$. By letting $\lambda = -\ln \kappa$, we can write $S(t) = e^{-\lambda t}$.

Problem 13. The gamma function is a generalized factorial function.

(a) Show $\Gamma\left(\frac{1}{2}\right) = 2 \int_0^{\infty} e^{-x^2} dx$.

Hint: Use the substitution $u = x^2$.

(b) Show $\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}$.

Hint: Use the result of optional problem 11.

(c) Show that $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$, and that

$$\Gamma\left(\frac{n}{2}\right) = \frac{\sqrt{\pi}(n-1)!}{2^{n-1} \left(\frac{n-1}{2}\right)!}$$

if n is an odd integer.

Answers to exercises

Problem 1. Let X be the life time of the electronic component.

(a) $P(X \leq 10) = 1 - e^{-(0.2)(10)} \approx 0.8647$

(b) $P(5 \leq X \leq 15) = e^{-(0.2)(5)} - e^{-(0.2)(15)} \approx 0.3181$

Problem 2. Let X be a normal random variable with $\mu = 70$ and $\sigma = 3$.

(a) $P(X \geq (6)(12)) = 1 - \Phi\left(\frac{72 - 70}{3}\right) \approx 1 - \Phi(0.67) \approx 0.25$

(b) In centimeters, the height becomes $(2.54)X$, and it is normally distributed with $\mu = (2.54)(70) = 177.8$ and $\sigma = (2.54)(3) = 7.62$.

Problem 3.

(a) $P(X > 10) = 1 - \Phi\left(\frac{10 - 5}{10}\right) = 1 - \Phi(0.5) = 0.3085$.

(b) $P(-20 < X < 15) = \Phi\left(\frac{15 - 5}{10}\right) - \Phi\left(\frac{-20 - 5}{10}\right) = \Phi(1) - \Phi(-2.5) = 0.8351$.

(c) Since $P(X > x) = 1 - \Phi\left(\frac{x - 5}{10}\right) = 0.05$, we must have $\Phi\left(\frac{x - 5}{10}\right) = 0.95$. By using the normal distribution table, we can find $\frac{x - 5}{10} \approx 1.64$. Thus, $x \approx 5 + (10)(1.64) = 21.4$.

Problem 4.

(a) $P(|X - \mu| \leq 0.675\sigma) = P(-0.675\sigma \leq X - \mu \leq 0.675\sigma) = \Phi(0.675) - \Phi(-0.675) \approx 0.5$

(b) $P(\mu - c \leq X \leq \mu + c) = 0.95$ implies $P\left(\frac{X - \mu}{\sigma} \leq \frac{c}{\sigma}\right) = \Phi\left(\frac{c}{\sigma}\right) = 0.975$. Then we can find $\frac{c}{\sigma} \approx 1.96$. Thus, $c = 1.96\sigma$.

Problem 5. By solving $P(X < 1) = 1 - e^{-\lambda} = 0.05$, we obtain $\lambda = -\ln 0.95 \approx 0.0513$.

Problem 6. (a) Let p_1 be the probability that the component breaks down before 5 months, and let p_2 be the probability that the component breaks down before 50 months. Then,

$$p_1 = 1 - e^{-5\mu} \approx 1 - (1 - 0.0005) = 0.0005;$$

$$p_2 = 1 - e^{-50\mu} \approx 1 - (1 - 0.005) = 0.005.$$

(b) Let X_1 be the number of components breaking down before 5 months, and let X_2 be the number of components breaking down before 50 months. Then X_i has the binomial distribution

with parameter $(p_i, 1000)$ for $i = 1, 2$. By using the Poisson approximation with $\lambda_1 = 1000p_1 = 0.5$ and $\lambda_2 = 1000p_2 = 5$, we can obtain

$$P(X_1 \leq 2) = e^{-\lambda_1} \left(1 + \lambda_1 + \frac{\lambda_1^2}{2} \right) \approx 0.986;$$

$$P(X_2 \leq 2) = e^{-\lambda_2} \left(1 + \lambda_2 + \frac{\lambda_2^2}{2} \right) \approx 0.125.$$

Problem 7. Let X be the length of phone call in minutes.

(a) By solving $P(X \leq 20) = 1 - e^{-20\lambda} = 0.5$, we obtain $\lambda = \frac{\ln 0.5}{20} \approx 0.0347$.

(b) $P(10 \leq X \leq 20) = (1 - e^{-20\lambda}) - (1 - e^{-10\lambda}) \approx 0.207$.

Problem 8. Since $Y = X_1^2 + X_2^2 + X_3^2 + X_4^2 + X_5^2 + X_6^2$ has a chi-square random variables with $n = 6$, we obtain $P(Y \leq 1.64) = 0.05$.

Problem 9. (a) Recall that $a < 0$.

$$\begin{aligned} P(Y \leq t) &= P(aX + b \leq t) = P(X \geq (t - b)/a) = \int_{(t-b)/a}^{\infty} \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{(x - \mu)^2}{2\sigma^2} \right] dx \\ &= \int_{-\infty}^t \frac{1}{|a|\sigma\sqrt{2\pi}} \exp \left[-\frac{(y - a\mu - b)^2}{2(a\sigma)^2} \right] dy \end{aligned}$$

by substituting $x = (y - b)/a$.

(b) The integrand $f(y)$ is a normal density with $(a\mu + b, (a\sigma)^2)$.

Problem 10. For $t > 0$ we obtain

$$\begin{aligned} P(Y \leq t) &= P(e^X \leq t) = P(X \leq \ln t) = \int_{-\infty}^{\ln t} \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{(x - \mu)^2}{2\sigma^2} \right] dx \\ &= \int_0^t \frac{1}{y\sigma\sqrt{2\pi}} \exp \left[-\frac{(\ln y - \mu)^2}{2\sigma^2} \right] dy \end{aligned}$$

by substituting $x = \ln y$. Thus, the density is given by

$$f(y) = \frac{1}{y\sigma\sqrt{2\pi}} \exp \left[-\frac{(\ln y - \mu)^2}{2\sigma^2} \right], \quad y > 0.$$