

Limiting Distributions

We introduce the mode of convergence for a sequence of random variables, and discuss the convergence in probability and in distribution. The concept of convergence leads us to the two fundamental results of probability theory: Law of large number and central limit theorem.

Indicator function. We can define an *indicator function* I_A of a subset A of real line as

$$I_A(x) = \begin{cases} 1 & x \in A; \\ 0 & \text{otherwise.} \end{cases}$$

Let X be a random variable having the pdf $f(x)$. Then we can express the probability

$$P(X \in A) = \int_A f(x) dx = \int_{-\infty}^{\infty} I_A(x) f(x) dx$$

Chebyshev's inequality. Suppose that X has the mean $\mu = E[X]$ and the variance $\sigma^2 = \text{Var}(X)$, and that we choose the particular subset

$$A = \{x : |x - \mu| > t\}.$$

Then we can observe that

$$I_A(x) \leq \frac{1}{t^2}(x - \mu)^2 \quad \text{for all } x,$$

which can be used to derive

$$\begin{aligned} P(|X - \mu| > t) &= P(X \in A) = \int_{-\infty}^{\infty} I_A(x) f(x) dx \\ &\leq \frac{1}{t^2} \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx = \frac{1}{t^2} \text{Var}(X) = \frac{\sigma^2}{t^2}. \end{aligned}$$

The inequality $P(|X - \mu| > t) \leq \sigma^2/t^2$ is called the *Chebyshev's inequality*.

Convergence in probability. Let Y_1, Y_2, \dots be a sequence of random variables. Then we can define the event $A_n = \{|Y_n - \alpha| > \varepsilon\}$ and the probability $P(A_n) = P(|Y_n - \alpha| > \varepsilon)$. We say that the sequence Y_n converges to α in probability if we have

$$\lim_{n \rightarrow \infty} P(|Y_n - \alpha| > \varepsilon) = 0,$$

for any choice of $\varepsilon > 0$. In general, the convergent value can be a random variable Y , and we say that Y_n converges to Y in probability if it satisfies

$$\lim_{n \rightarrow \infty} P(|Y_n - Y| > \varepsilon) = 0,$$

for any choice of $\varepsilon > 0$.

Laws of large numbers. Now let X_1, X_2, \dots be a sequence of iid random variables with mean μ and variance σ^2 . Then for each $n = 1, 2, \dots$, the sample mean

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

has the mean μ and the variance σ^2/n . Given a fixed $\varepsilon > 0$, we can apply the Chebyshev's inequality to obtain

$$P(|\bar{X}_n - \mu| > \varepsilon) \leq \frac{\sigma^2/n}{\varepsilon^2} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Laws of large numbers, continued. We have established the assertion

“the sample mean \bar{X}_n converges to μ in probability”

which is called the *weak law of large numbers*. A stronger result of the above statement, “the sample mean \bar{X}_n converges to μ almost surely,” was established by Kolmogorov, and is called the *strong law of large numbers*.

Convergence in distribution. Let X_1, X_2, \dots be a sequence of random variables having the cdf's F_1, F_2, \dots , and let X be a random variable having the cdf F . Then we say that the sequence X_n converges to X in distribution (in short, X_n converges to F) if

$$\lim_{n \rightarrow \infty} F_n(x) = F(x) \quad \text{for every } x \text{ at which } F(x) \text{ is continuous.} \quad (7.1)$$

Convergence in distribution, continued. Convergence in distribution does not necessarily imply convergence in probability.

Example 1. Let Y be a uniform random on $[-1/2, 1/2]$, and let $X_1 = Y, X_2 = -Y, X_3 = Y, X_4 = -Y, \dots$. Then we have

$$F_n(t) = F_Y(t) = \begin{cases} 0 & \text{if } t < -1/2; \\ t + 1/2 & \text{if } -1/2 \leq t \leq 1/2; \\ 1 & \text{if } t > 1/2. \end{cases}$$

Then clearly X_n converges to Y in distribution. Does X_n converges to Y in probability?

Solution to Example 1. Let $0 < \varepsilon < 1$ be arbitrary. Since we have

$$P(|X_n - Y| > \varepsilon) = \begin{cases} 0 & \text{if } n \text{ is odd;} \\ P(|2Y| > \varepsilon) = 1 - \varepsilon & \text{if } n \text{ is even,} \end{cases}$$

$\lim_{n \rightarrow \infty} P(|X_n - Y| > \varepsilon)$ does not exist. Thus, X_n does not converge to Y in probability. In fact, the limit in probability does not exist.

Convergence in distribution, continued. Convergence in distribution does not necessarily establish $\lim_{n \rightarrow \infty} F_n(x) = F(x)$ for every point.

Example 2. Let X_n be a uniform random on $[-1/n, 1/n]$, and let $X \equiv 0$ be the constant.

$$F_n(t) = \begin{cases} 0 & \text{if } t < -1/n; \\ (nt + 1)/2 & \text{if } -1/n \leq t \leq 1/n; \\ 1 & \text{if } t > 1/n. \end{cases}$$

Does X_n converges to X in distribution? In probability?

Solution to Example 2. Observe that the constant $X \equiv 0$ has the cdf

$$F(t) = \begin{cases} 0 & \text{if } t < 0; \\ 1 & \text{if } t \geq 0. \end{cases}$$

If $t < 0$ then $\lim_{n \rightarrow \infty} F_n(t) = 0 = F(t)$; if $t > 0$ then $\lim_{n \rightarrow \infty} F_n(t) = 1 = F(t)$. However, we can observe

$$\lim_{n \rightarrow \infty} F_n(0) = 1/2 \neq F(0).$$

Since $F(t)$ is not continuous at $t = 0$, X_n converges to X in distribution. Since for any $\varepsilon > 0$ we have

$$\lim_{n \rightarrow \infty} P(|X_n - X| > \varepsilon) = \lim_{n \rightarrow \infty} P(|X_n| > \varepsilon) = 0,$$

it converges in probability.

Alternative criterion for convergence in distribution. Let Y_1, Y_2, \dots be a sequence of random variables, and let Y be a random variable. Then the sequence Y_n converges to Y in distribution if and only if

$$\lim_{n \rightarrow \infty} E[g(Y_n)] = E[g(Y)]$$

for every continuous function g satisfying $E[|g(Y_n)|] < \infty$ for all n . By setting $g(y) = e^{ty}$, we obtain the moment generating function $M_Y(t) = E[g(Y)]$. Thus, if Y_n converges to Y in distribution then

$$\lim_{n \rightarrow \infty} M_{Y_n}(t) = M_Y(t)$$

when $M_{Y_n}(t)$ exists for all n .

Alternative criterion, continued. Convergence in (7.1) is completely characterized in terms of the distributions F_1, F_2, \dots and F . Recall that the distributions $F_1(x), F_2(x), \dots$ and $F(x)$ are uniquely determined by the respective moment generating functions, say $M_1(t), M_2(t), \dots$ and $M(t)$. Furthermore, we have an “equivalent” version of the convergence in terms of the m.g.f.’s.

Theorem 3. *If*

$$\lim_{n \rightarrow \infty} M_n(t) = M(t) \quad \text{for every } t \text{ (around } 0), \quad (7.2)$$

then the corresponding distributions F_n ’s converge to F in the sense of (7.1).

Slutsky’s theorem. Although the following theorem is technical, it plays a critical role later in asymptotic theory of statistics.

Theorem 4. *Let Z_1, Z_2, \dots and W_1, W_2, \dots be two sequences of random variables, and let c be a constant value. Suppose that the sequence Z_n converges to Z in distribution, and that the sequence W_n converges to c in probability. Then*

(a) *The sequence $Z_n + W_n$ converges to $Z + c$ in distribution.*

(b) *The sequence $Z_n W_n$ converges to cZ in distribution.*

(c) *Assuming $c \neq 0$, the sequence Z_n/W_n converges to Z/c in distribution.*

Central limit theorem. Central limit theorem (CLT) is the most important theorem in probability and statistics.

Theorem 5. Let X_1, X_2, \dots be a sequence of iid random variables having the common distribution F with mean 0 and variance 1. Then

$$Z_n = \sum_{i=1}^n X_i / \sqrt{n} \quad n = 1, 2, \dots$$

converges to the standard normal distribution.

A sketch of proof. Let M be the m.g.f. for the distribution F , and let M_n be the m.g.f. for the random variable Z_n for $n = 1, 2, \dots$. Since X_i 's are iid random variables, we have

$$\begin{aligned} M_n(t) &= E[e^{tZ_n}] = E\left[e^{\left(\frac{t}{\sqrt{n}}\right)(X_1 + \dots + X_n)}\right] \\ &= E\left[e^{\left(\frac{t}{\sqrt{n}}\right)X_1}\right] \times \dots \times E\left[e^{\left(\frac{t}{\sqrt{n}}\right)X_n}\right] = \left[M\left(\frac{t}{\sqrt{n}}\right)\right]^n \end{aligned}$$

A sketch of proof, continued. Observing that $M'(0) = E[X_1] = 0$ and $M''(0) = E[X_1^2] = 1$, we can apply a Taylor expansion to $M\left(\frac{t}{\sqrt{n}}\right)$ around 0.

$$\begin{aligned} M\left(\frac{t}{\sqrt{n}}\right) &= M(0) + \frac{t}{\sqrt{n}}M'(0) + \frac{t^2}{2n}M''(0) + \varepsilon_n(t) \\ &= 1 + \frac{t^2}{2n} + \varepsilon_n(t). \end{aligned}$$

Recall that $\lim_{n \rightarrow \infty} \left(1 + \frac{a}{n}\right)^n = e^a$; in a similar manner, we can obtain

$$\lim_{n \rightarrow \infty} M_n(t) = \lim_{n \rightarrow \infty} \left[1 + \frac{t^2/2}{n} + \varepsilon_n(t)\right]^n = e^{t^2/2},$$

which is the m.g.f. for the standard normal distribution.

Central limit theorem, modified. Mean and variance for a sequence can be arbitrary in general, and CLT can be modified for them.

Theorem 6. Let X_1, X_2, \dots be a sequence of iid random variables having the common distribution F with mean μ and variance σ^2 . Then

$$Z_n = \frac{\sum_{i=1}^n X_i - n\mu}{\sigma\sqrt{n}} = \frac{\sum_{i=1}^n (X_i - \mu)/\sigma}{\sqrt{n}} \quad n = 1, 2, \dots$$

converges to the standard normal distribution.

Normal approximation. Let X_1, X_2, \dots, X_n be finitely many iid random variables with mean μ and variance σ^2 . If the size n is adequately large, then the distribution of the sum

$$Y = \sum_{i=1}^n X_i$$

can be approximated by the normal distribution with parameter $(n\mu, n\sigma^2)$. A general rule for “adequately large” n is about $n \geq 30$, but it is often good for much smaller n .

Normal approximation, continued. Similarly the sample mean

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

has approximately the normal distribution with parameter $(\mu, \sigma^2/n)$. Then the following application of normal approximation to probability computation can be applied.

Random variable	Approximation	Probability
$Y = X_1 + \dots + X_n$	$N(n\mu, n\sigma^2)$	$\Phi\left(\frac{b-n\mu}{\sigma\sqrt{n}}\right) - \Phi\left(\frac{a-n\mu}{\sigma\sqrt{n}}\right)$
$\bar{X} = \frac{X_1 + \dots + X_n}{n}$	$N\left(\mu, \frac{\sigma^2}{n}\right)$	$\Phi\left(\frac{b-\mu}{\sigma/\sqrt{n}}\right) - \Phi\left(\frac{a-\mu}{\sigma/\sqrt{n}}\right)$

Example 7. An insurance company has 10,000 automobile policyholders. The expected yearly claim per policyholder is \$240 with a standard deviation of \$800. Find approximately the probability that the total yearly claim exceeds \$2.7 million. Can you say that such event is statistically highly unlikely?

Solution. Let X_1, \dots, X_{10000} be normally distributed with mean $\mu = 240$ and $\sigma = 800$. Then $Y = X_1 + \dots + X_{10000}$ is approximately normally distributed with mean $n\mu = 2,400,000$ and standard deviation $\sqrt{n}\sigma = 80,000$. Thus, we obtain

$$P(Y > 2,700,000) \approx 1 - \Phi(3.75) \approx 0,$$

which is highly unlikely.

Normal approximation to binomial distribution. Suppose that X_1, \dots, X_n are iid Bernoulli random variables with the mean $p = E(X)$ and the variance $p(1 - p) = \text{Var}(X)$. If the size n is adequately large, then the distribution of the sum

$$Y = \sum_{i=1}^n X_i$$

can be approximated by the normal distribution with parameter $(np, np(1 - p))$. Thus, the normal distribution $N(np, np(1 - p))$ approximates the binomial distribution $B(n, p)$. A general rule for “adequately large” n is to satisfy $np \geq 5$ and $n(1 - p) \geq 5$.

Normal approximation to binomial distribution, continued. Let Y be a binomial random variable with parameter (n, p) , and let Z be a normal random variable with parameter $(np, np(1-p))$. Then the distribution of Y can be approximated by that of Z . Since Z is a continuous random variable, the approximation of probability should improve when the following formula of *continuity correction* is considered.

$$\begin{aligned} P(i \leq Y \leq j) &\approx P(i - 0.5 \leq Z \leq j + 0.5) \\ &= \Phi\left(\frac{j + 0.5 - np}{\sqrt{np(1-p)}}\right) - \Phi\left(\frac{i - 0.5 - np}{\sqrt{np(1-p)}}\right). \end{aligned}$$

Exercises

Problem 1. Let S_n^2 denote the sample variance of random sample of size n from normal distribution with mean μ and variance σ^2 .

- (a) Find $E[S_n^2]$ and $\text{Var}(S_n^2)$.
- (b) Show that S_n^2 converges to σ^2 in probability

Problem 2. Let X_1, \dots, X_n be a random sample of size n from the common pdf $f(x) = e^{-(x-\theta)}$ if $\theta \leq x < \infty$; otherwise, $f(x) = 0$, and let $X_{(1)}$ be the first order statistic, and $Z_n = n(X_{(1)} - \theta)$.

- (a) Find the pdf for $X_{(1)}$, and the mgf $M(t)$.
- (b) Find the mgf $M_n(t)$ for Z_n .
- (c) Does Z_n converge in distribution? If so, find the limiting distribution.

Problem 3. Suppose that Z_n has a Poisson distribution with parameter $\lambda = n$. Then show that $(Z_n - n)/\sqrt{n}$ converges to the standard normal distribution.

Problem 4. An actual voltage of new a 1.5-volt battery has the probability density function

$$f(x) = 5, \quad 1.4 \leq x \leq 1.6.$$

Estimate the probability that the sum of the voltages from 120 new batteries lies between 170 and 190 volts.

Problem 5. The germination time in days of a newly planted seed has the probability density function

$$f(x) = 0.3e^{-0.3x}, \quad x \geq 0.$$

If the germination times of different seeds are independent of one another, estimate the probability that the average germination time of 2000 seeds is between 3.1 and 3.4 days.

Problem 6. Calculate the following probabilities by using normal approximation with continuity correction.

- (a) Let X be a binomial random variable with $n = 10$ and $p = 0.7$. Find $P(X \geq 8)$.
- (b) Let X be a binomial random variable with $n = 15$ and $p = 0.3$. Find $P(2 \leq X \leq 7)$.
- (c) Let X be a binomial random variable with $n = 9$ and $p = 0.4$. Find $P(X \leq 4)$.
- (d) Let X be a binomial random variable with $n = 14$ and $p = 0.6$. Find $P(8 \leq X \leq 11)$.

Optional problems

Convergence in distribution.

Definition 8. A function $f(x)$ is said to be *bounded* if there exists some $M > 0$ such that $|f(x)| \leq M$ for all x . A function $f(x)$ is called *uniformly continuous* if for any $\varepsilon > 0$ there exists some $\delta > 0$ such that $|f(x) - f(y)| < \varepsilon$ whenever $|x - y| < \delta$.

Let X_1, X_2, \dots be a sequence of random variables, and let X be a random variable. In order to see that X_n converges to X in distribution it suffices to show that

$$\lim_{n \rightarrow \infty} E[g(X_n)] = E[g(X)]$$

for every bounded and uniformly continuous function f .

Problem 7. Choose a bounded and uniformly continuous function f so that $|f(x)| \leq M$ for all x and $|f(x) - f(y)| < \varepsilon$ whenever $|x - y| < \delta$. Let X be a random variable, and let $A_\delta = \{y : |y - c| < \delta\}$. Suppose that a sequence Y_n of random variables converges to a constant c in probability.

- (a) Show that $|f(X + Y_n) - f(X + c)| \leq 2M$.
- (b) Show that $|f(X + Y_n) - f(X + c)|I_{A_\delta}(Y_n) \leq \varepsilon$.
- (c) Show that $|E[f(X + Y_n)] - E[f(X + c)]| \leq 2MP(|Y_n - c| \geq \delta) + \varepsilon$
- (d) The above result implies that $\lim_{n \rightarrow \infty} |E[f(X + Y_n)] - E[f(X + c)]| \leq \varepsilon$ for every $\varepsilon > 0$, and therefore, that $\lim_{n \rightarrow \infty} |E[f(X + Y_n)] - E[f(X + c)]| = 0$. Then argue that $X + Y_n$ converges to $X + c$ in distribution.

Problem 8. Let Z_1, Z_2, \dots and W_1, W_2, \dots be two sequences of random variables, and let c be a constant value. Suppose that Z_n converges to Z in distribution, and that W_n converges to c in probability.

- (a) Choose a bounded and uniformly continuous function f so that $|f(x)| \leq M$ for all x and $|f(x) - f(y)| < \varepsilon$ whenever $|x - y| < \delta$. Then show that

$$\begin{aligned} & |E[f(Z_n + W_n)] - E[f(Z + c)]| \\ & \leq |E[f(Z_n + c)] - E[f(Z + c)]| + |E[f(Z_n + W_n)] - E[f(Z_n + c)]| \\ & \leq |E[f(Z_n + c)] - E[f(Z + c)]| + 2MP(|W_n - c| \geq \delta) + \varepsilon \end{aligned}$$

- (b) The above result implies that $\lim_{n \rightarrow \infty} |E[f(Z_n + W_n)] - E[f(Z + c)]| \leq \varepsilon$, and therefore, that $\lim_{n \rightarrow \infty} |E[f(Z_n + W_n)] - E[f(Z + c)]| = 0$. Then argue that $Z_n + W_n$ converges to $Z + c$ in distribution (Slutsky's theorem).

Answers to exercises

Problem 1. Recall that $W_{n-1} = (n-1)S_n^2/\sigma^2$ is a chi-square distribution with $(n-1)$ degrees of freedom, and that $E[W_{n-1}] = (n-1)$ and $\text{Var}(W_{n-1}) = 2(n-1)$.

(1) $E[S_n^2] = E\left[\frac{\sigma^2 W_{n-1}}{(n-1)}\right] = \frac{\sigma^2}{n-1}E[W_{n-1}] = \sigma^2$, and

$$\text{Var}(S_n^2) = \text{Var}\left(\frac{\sigma^2 W_{n-1}}{(n-1)}\right) = \left(\frac{\sigma^2}{n-1}\right)^2 \text{Var}(W_{n-1}) = \frac{2\sigma^4}{n-1}.$$

(2) By Chebyshev's inequality we obtain

$$P(|S_n^2 - \sigma^2| > \varepsilon) \leq \frac{2\sigma^4}{(n-1)\varepsilon^2} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Problem 2. (1) $X_{(1)}$ has the pdf $f_{(1)}(x) = ne^{-n(x-\theta)}$ if $x \geq \theta$; otherwise, $f_{(1)}(x) = 0$. Then we obtain

$$M(t) = \int_{\theta}^{\infty} e^{tx} ne^{-n(x-\theta)} dx = ne^{t\theta} \int_0^{\infty} e^{-(n-t)u} du = \frac{ne^{t\theta}}{n-t}$$

for $t < n$.

(2) We obtain

$$M_n(t) = E[e^{tn(X_{(1)}-\theta)}] = e^{nt\theta} M(nt) = \frac{1}{1-t}$$

for $t < 1$.

(3) Since $\lim_{n \rightarrow \infty} M_n(t) = \frac{1}{1-t}$, the limiting distribution is an exponential with $\lambda = 1$.

Problem 3. Let X_1, X_2, \dots be iid random variables having a Poisson distribution with $\lambda = 1$. Then we have $Z_n = X_1 + \dots + X_n$. By CLT we obtain

$$Z_n = \frac{\sum_{i=1}^n X_i - n}{\sqrt{n}}$$

converges to the standard normal distribution.

Problem 4. Let X_1, \dots, X_{120} be the voltage of each battery, having the pdf $f(x)$. Since $\mu = E[X_i] = 1.5$ and $\sigma^2 = E[X_i^2] - \mu^2 = 0.0033$, the sum $Y = \sum_{i=1}^{120} X_i$ is approximately distributed as $N(120\mu, 120\sigma^2) = N(180, 0.396)$. Thus, we obtain

$$P(170 \leq Y \leq 190) = \Phi\left(\frac{190 - 180}{\sqrt{0.396}}\right) - \Phi\left(\frac{170 - 180}{\sqrt{0.396}}\right) \approx 1$$

Problem 5. Let X_1, \dots, X_{2000} be the germination time of individual seed, having the exponential distribution with $\lambda = 0.3$. Since $\mu = E[X_i] = 1/0.3 \approx 3.33$ and $\sigma^2 = \text{Var}(X_i) = 1/0.3^2 \approx 11.11$, the sample mean \bar{X} is approximately distributed as $N(\mu, \sigma^2/n) = N(3.33, 0.0056)$. Thus,

$$P(3.1 \leq \bar{X} \leq 3.4) = \Phi\left(\frac{3.4 - 3.33}{\sqrt{0.0056}}\right) - \Phi\left(\frac{3.1 - 3.33}{\sqrt{0.0056}}\right) \approx 0.825$$

Problem 6. (1) $P(X \geq 8) = 1 - \Phi\left(\frac{7.5 - (10)(0.7)}{\sqrt{(10)(0.7)(0.3)}}\right) \approx 0.365$

(2) $P(2 \leq X \leq 7) = \Phi\left(\frac{7.5 - (15)(0.3)}{\sqrt{(15)(0.3)(0.7)}}\right) - \Phi\left(\frac{1.5 - (15)(0.3)}{\sqrt{(15)(0.3)(0.7)}}\right) \approx 0.909$

(3) $P(X \leq 4) = \Phi\left(\frac{4.5 - (9)(0.4)}{\sqrt{(9)(0.4)(0.6)}}\right) \approx 0.7299$

(4) $P(8 \leq X \leq 11) = \Phi\left(\frac{11.5 - (14)(0.6)}{\sqrt{(14)(0.6)(0.4)}}\right) - \Phi\left(\frac{7.5 - (14)(0.6)}{\sqrt{(14)(0.6)(0.4)}}\right) \approx 0.6429$